

# The transcriptome of acute dehydration in myeloid leukemic cells

## ABSTRACT

Human myeloid leukemia cells (HL-60/S4) exposed to hyperosmotic stress with sucrose undergo dehydration and cell shrinkage. Interphase chromatin and mitotic chromosomes congeal, exhibiting altered phase separation (demixing) of chromatin proteins. To investigate changes in the transcriptome, we exposed HL-60/S4 cells to hyperosmotic sucrose stress (~600 milliOsmolar) for 30 and 60 minutes. We employed RNA-Seq of polyA mRNA to identify genes with increased or decreased transcript levels relative to untreated control cells (i.e., differential gene expression). These genes were examined for over-representation of Gene Ontology (GO) terms. In stressed cells, multiple GO terms associated with transcription, translation, mitochondrial function and proteasome activity, as well as “replication-dependent histones”, were over-represented among genes with increased transcript levels; whereas, genes with decreased transcript levels were over-represented with transcription repressors. The transcriptome profiles of hyperosmotically-stressed cells suggest acquisition of cellular rebuilding, a futile homeostatic response, as these cells are ultimately doomed to a dehydrated death.

## INTRODUCTION

In a beautiful and convincing study of buffer conditions used for the isolation of mammalian interphase nuclei [1], the authors demonstrated that employing uncharged, inert, water-soluble polymers in the nuclear isolation buffer preserved ultrastructure and transcription capability. The polymers cannot traverse an intact nuclear envelope; they simulate the cytoplasmic iso-osmotic pressure on the nucleus. Normal cells in our body are frequently exposed to hyperosmotic stress conditions [2]. These are generally brief challenges of resilient cell systems. More serious consequences can occur during tissue inflammation and various diseases [2]. There has been considerable exploration of the adaptation of renal (kidney) cells to hyperosmotic stress. However, evidence suggests that other tissues respond differently [3] and there is a need to develop convenient reproducible models using other cell types. The influence of growth medium osmolarity upon *in vivo* mammalian tissue culture cell nuclear structure has been explored for several decades. Despite the variation of cell types studied, there is a consensus that acute hyperosmotic stress (e.g., >300 milliOsmolar [mOsM], total) produces rapid cell volume shrinkage and heterogeneity of nuclear chromatin condensation [4-8]. In our previous study [8], we exposed live human leukemic HL-60/S4 cells to 300 mM sucrose in growth medium (i.e., ~600 mOsM, total) for 30 and 60 minutes, producing acute dehydration and cell shrinkage. Employing microscopy, we observed interphase and mitotic chromatin condensation (denoted by us: “congelation” \*\*). The effect on interphase nuclear chromatin can be clearly seen employing DAPI staining of DNA combined with STED imaging (Fig. 1): the dispersed “fine” interphase nuclear chromatin fibers (in the absence of sucrose) “congeal” into thicker strands during hyperosmotic stress. Combined with immunofluorescent staining microscopy, we documented apparent phase separation with loss of colocalization (possible “demixing”) of various chromatin-associated proteins resulting from the acute hyperosmotic stress [8]. In the present study, we do not focus on long-term incubation (i.e., >1 hour) of HL-60/S4 cells in 300 mM sucrose, since the cells appear to deteriorate and die after several hours in these dehydrating conditions. In this regard, HL-60/S4 cells appear to differ from some other cell types that can adapt to ongoing hyperosmotic stress [9,10].

Two central questions of the present study are: 1) How is the polyA mRNA transcriptome of cells exposed to acute hyperosmotic stress altered, in comparison to unstressed cells? 2) How can this altered transcriptome be interpreted

\*\*Footnote: The chromatin description “Congelation” was first employed in [8]. This chosen description is intended to distinguish hyperosmotic stress chromatin condensation from normal heterochromatin or mitotic chromosome condensation.

David B. Mark Welch<sup>1</sup>,

Travis J. Gould<sup>2</sup>,

Ada L. Olins<sup>3</sup>,

Donald E. Olins<sup>3</sup>✉

<sup>1</sup>Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole MA, USA

<sup>2</sup>Department of Physics & Astronomy, Bates College, Lewiston, ME, USA

<sup>3</sup>MaineHealth Institute for Research (MHIR), Scarborough, ME USA

[https://doi.org/10.18388/pb.2021\\_507](https://doi.org/10.18388/pb.2021_507)

✉Corresponding Author: [donaolins@gmail.com](mailto:donaolins@gmail.com)

## DEDICATION

This article is dedicated to the memory of Ronald Hancock, who's excellent studies on “nuclear crowding” inspired the authors to explore the effects of hyperosmotic stress on nuclear chromatin structure and mRNA transcription. Ron was a fine scientist and a generous human being. We would have liked to hear his reaction to this study.

**Keywords:** Cellular dehydration, RNA polymerase, Ribosomal proteins, Mitochondrial Function, Histones

**Abbreviations:** DGE – differential gene expression; ORA – over-representation-analysis; GO – gene ontology; HGCN – HUGO Gene Nomenclature Committee

**Table 1.** Number of genes with significant changes in relative transcript levels for HL-60/S4 differentiation states [13] and for 30- and 60-minute exposure times to 300 mM sucrose (present study).

DGE	Differentiation				Sucrose			
	RA	%	TPA	%	30 min	%	60 min	%
Increased	4,249	26.2	5,156	31.2	3,128	18.4	3,231	20.3
Decreased	3,900	24.1	4,528	27.4	2,746	20.9	2,962	22.2
Unchanged	8,040	49.7	6,842	41.4	9,073	60.6	8,371	57.5
Total	16,189		16,526		14,947		14,564	
Control	15,998				15,615			

Column titles: DGE, Differential Gene Expression; RA, 4 days of phorbol ester differentiation to macrophage form; 30 and 60 min, exposure time in medium+300 mM sucrose. "Control" is the total number of genes mapped under control conditions. "%" is the percentage of the appropriate "Total" for the specific cell condition. For a complete list of all analyzed 25,351 genes, see Table S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>.

to understand changes in the stressed cell physiological state? Two prior studies have been published toward this end; both employed microarray analyses of several hundred genes [11,12]. Because these studies employed a different analysis method and examined different cell types than to the present study, a direct comparison is difficult. In the present study, we compared the transcriptomes of acute (i.e., 30 and 60 min.) hyperosmotically-stressed undifferentiated HL-60/S4 cells to that of unstressed (control) cells. HL-60/S4 cells are derived from a human myeloid leukemia that preserves the ability to differentiate *in vitro* into stable cell states [13,14]. We employed a standard RNA-Seq approach using polyA mRNA isolated from biological replicates of stressed and unstressed cells, and mapped reads back to the human genome as a proxy for the equilibrium mRNA transcript level of each gene at the time of sampling. Differential gene expression (DGE) analysis of ~16,000 genes revealed statistically significant increases or decreases in transcript levels compared to control (unstressed) transcript levels. We then employed over-representation-analysis (ORA) to identify gene ontology (GO) terms enriched in genes with increased or decreased transcript levels. The resulting profiles of GO terms permit us to speculate on the probable physiological functions that are affected in the stressed cells (i.e., the altered cell physiological state).


## RESULTS

### OVERALL VIEW OF ACUTE HYPEROSMOTIC STRESS EFFECTS ON THE mRNA TRANSCRIPTOME

Despite the considerable extent of interphase chromatin congelation at 30 and 60 minutes of hyperosmotic sucrose treatment, the relative transcript levels are essentially unchanged for most (~60%) genes (Table 1 "Sucrose"). After 30 minutes of exposure, transcript levels are significantly increased for 3128 genes and significantly decreased for 2746. This change in transcript levels continues with ongoing exposure to sucrose: between 30 and 60 minutes the number of genes with significantly increased or decreased transcript levels changed by 103 and 215 genes, respectively. See Table S1, at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for EBseq normalized mean counts and PPDE (statistical significance) values for each gene at a pairwise comparison of conditions. The implication of these results is that most of the transcriptome changes occurred within the initial 30 minutes. It is of interest to compare these hyper-

osmotic stress transcriptome changes to the transcriptome changes of chemically induced differentiation in HL-60/S4 cells; i.e., 4 days with retinoic acid (RA) into granulocytes or 4 days with phorbol ester (TPA) into macrophage [13]. The differentiated cell states only exhibited a slightly greater change in relative transcript levels (Table 1 "Differentiation"). Observing cell differentiation in the microscope indicated that during the 4 day induction period visible cellular phenotypes exhibited considerable morphological changes (Fig. 1 of [14] and Fig. 5 of [13]). It is important to point-out that the cells tested by hyperosmotic stress (this study) were undifferentiated. A lower percentage of genes remained unchanged after 4 days of differentiation, compared to the percentage of unchanged genes after 30 and 60 minutes of hyperosmotic sucrose stress.

We performed over-representation-analysis (ORA) of genes with increased and decreased transcript levels using each of the three GO non-redundant annotations (i.e., Biological Processes, Cellular Component and Molecular Function). Analysis of genes with increased transcript levels (upregulation) revealed GO terms associated with transcription and translation, mitochondrial structure and function, and protein stress (Table 2; also, see Tables S2, S3 and S4 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for complete ORA results with GO term identifiers, enrichment, and significance). The most enriched GO



**Figure 1.** Effect of acute hyperosmotic stress upon interphase nuclear DNA distribution. STED microscopy images of undifferentiated HL-60/S4 cells after fixation, permeabilization and staining for DNA with DAPI. (Left Panel) Control cell nucleus, untreated in tissue culture medium. (Right Panel) Cell nucleus in tissue culture medium plus 300 mM sucrose for 30 minutes (~600 mOsm). Magnification bar: 10  $\mu$ m. Similar images can be found in Fig. 4 [8].

**Table 2.** Summary of GO terms from each functional domain which is enriched in genes with increased relative transcript levels in response to acute hyperosmotic stress, exposed for 0–30 min, 0–60 min and 30–60 min intervals.

GO terms associated with transcription and translation				GO terms associated with mitochondrial structure and function				GO terms associated with stress, proteasome, repair			
Biological Process	30	60	30-60	Cellular Component	30	60	30-60	Molecular Function	30	60	30-60
mRNA processing	•	•		spliceosomal complex	•	•	•	catalytic activity, acting on RNA	•	•	•
ribonucleoprotein complex biogenesis	•	•	•	Sm-like protein family complex	•	•	•	mRNA binding	•	•	•
ncRNA processing	•	•	•	small nucleolar ribonucleoprotein complex	•	•	•	snoRNA binding	•	•	•
RNA splicing	•	•	•	ribosome	•	•	•	structural constituent of ribosome	•	•	•
RNA catabolic process	•	•	•	preribosome	•	•	•	rRNA binding	•	•	•
ribonucleoprotein complex subunit organization	•	•	•	polysome	•	•	•	translation factor activity, RNA binding	•	•	•
RNA localization	•	•	•	nucleolar part	•	•	•	unfolded protein binding	•	•	•
rRNA metabolic process	•	•	•					ribonucleoprotein complex binding	•	•	•
tRNA metabolic process	•	•	•					tRNA binding	•	•	•
translational initiation	•	•	•					NF-kappaB binding	•	•	•
cytoplasmic translation	•	•	•								
translational elongation	•	•	•								
protein targeting	•	•	•	mitochondrial matrix	•	•	•	oxidoreductase activity, acting on NAD(P)H	•	•	•
nucleoside monophosphate metabolic process	•	•	•	mitochondrial inner membrane	•	•	•	electron transfer activity	•	•	•
establishment of protein localization to membrane	•	•	•	mitochondrial protein complex	•	•	•	oxidoreductase activity, acting on a heme group of donors	•	•	•
nucleoside triphosphate metabolic process	•	•	•	oxidoreductase complex	•	•	•				
mitochondrial gene expression	•	•	•	respiratory chain	•	•	•				
ribonucleotide metabolic process	•	•	•	NADH dehydrogenase complex	•	•	•				
mitochondrial respiratory chain complex assembly	•	•	•	organelle envelope lumen	•	•	•				
NADH dehydrogenase complex assembly	•	•	•	vesicle lumen	•	•	•				
generation of precursor metabolites and energy	•	•	•	cytochrome complex	•	•	•				
cytochrome complex assembly	•	•	•								
mitochondrial transport	•	•	•								
protein folding	•	•	•	peptidase complex	•	•	•	threonine peptidase activity	•	•	•
DNA damage response, detection of DNA damage	•	•	•	chaperone complex	•	•	•	chaperone binding	•	•	•
nucleotide-excision repair	•	•	•	primary lysosome	•	•	•	unfolded protein binding	•	•	•
proteasomal protein catabolic process	•	•	•	microbody	•	•	•	heat shock protein binding	•	•	•
protein-containing complex disassembly	•	•	•					oxidoreductase activity, acting on a sulfur group of donors	•	•	•
RNA catabolic process	•	•	•								
peroxisomal transport	•	•	•								
response to topologically incorrect protein	•	•	•								

• Significant using all genes with increased transcript levels      • Significant using genes with > 2-fold increase in transcript levels

Color coding: light green, enriched among genes with increased transcript levels, where  $\log_2FC > 0$ ; dark green, enriched among genes with increased transcript levels, where  $\log_2FC > 1$  (i.e., at least a doubling compared to control levels). See Tables S2, S3 and S4 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for complete GO data in the three annotations (S2 Biological Processes; S3 Cellular Component; S4 Molecular Function). These three Supporting Tables document the number of genes in each GO term that are upregulated and downregulated in 0–30 min, 0–60 min and 30–60 min intervals. For definitions of GO terms and lists of included genes, see <http://geneontology.org>

terms were associated with mitochondrial structure and function, which accounted for nearly 13% of all genes with increased transcript levels. In contrast, analysis of genes with decreased transcript levels yielded more diverse results, with GO terms reflecting a decrease in pathways involved in transcription repression and heterochromatin formation. Overall, these results suggest that the dehydrated physiological state of the stressed HL-60/S4 cells involves attempts to accomplish transcription and translation, and to increase mitochondrial oxidative phosphorylation, possibly a “rebuilding” phenotype to contend with protein misfolding and proteome degradation. Table 2 also shows the differences between 30 and 60 minutes of GO terms, demonstrating continued significant changes from 30 to 60 minutes for some GO terms.

#### NUCLEAR DISTRIBUTION OF “ACTIVATED” RNA POLYMERASE II

Observing the structural transition of dehydrated interphase nuclear chromatin (Fig. 1), we wondered how the spatial distribution of “activated” RNA Polymerase II (phosphorylated RNA Pol II) would be affected by hyperosmotic stress, relative to its distribution in untreated cells. Figure 2 displays confocal immunostaining images of anti-phosphorylated RNA Pol II, in relation to DAPI stained interphase nuclear chromatin ( $\pm 300$  mM sucrose for 30 minutes). These images clearly indicate that “activated” RNA


Pol II persists within the interphase nuclei of hyperosmotically stressed cells. However, there is an apparent redistribution of “activated” Pol II staining to locations near the surface of congealed chromatin and within spaces between congealed chromatin.

#### SELECTED EXAMPLES OF THE TRANSCRIPTOME DATA

Given the extent and depth of the mRNA transcriptome data, we have chosen to discuss only a few prominent stress-upregulated cellular functions. These data are presented in two types of graphs: 1) “Bar graphs”, showing the change in relative transcript levels (“Log2FC”,  $\log_2$  of the ratio of transcript levels) of genes representative for specific GO terms or other summaries of biological function. 2) “MA plots”, illustrating the relationship between Log2FC and mean transcript level for a population of related genes. In all cases, the transcriptome data is based upon all overlapping transcripts from a common promoter, rather than splice variants (i.e., entire genes rather than isoforms).

##### I. Transcription

The structure and function of eukaryotic RNA polymerases has been recently reviewed [17]. The three RNA polymerases (i.e., Pol I, Pol II, and Pol III, which synthesize primarily ribosomal RNA, protein coding mRNA, and tRNA, respectively) possess both specific and common pro-




**Figure 2.** Effect of hyperosmotic stress upon interphase distribution of “activated” phosphorylated RNA Polymerase II. Confocal images of undifferentiated HL-60/S4 cells after formaldehyde fixation ( $\pm 300$  mM sucrose), permeabilization and immunostaining. (Left Column) Control cells, untreated with sucrose in tissue culture medium. Note the absence of “activated” RNA Polymerase within mitotic chromosomes (yellow arrowhead), but presence within interphase nuclei. (Right Column) Interphase cell nuclei in tissue culture medium containing 300 mM sucrose ( $\sim 600$  mOsm) for 30 minutes prior to fixation. Staining: pPol II, phosphorylated (activated) RNA Polymerase II (green); PL2-6, anti-epichromatin, i.e., exposed nucleosome acidic patches at the surface of interphase chromatin (red), see references [8,15,16]; DAPI, DNA (blue). Magnification bar for all images: 10  $\mu$ m.

tein subunits. Our transcriptome data indicates that multiple subunits for each polymerase, including all of the common subunit genes increase their relative transcript levels during acute sucrose stress (Fig. 3 and Table S5 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>). Combined with Fig. 2, which establishes the persistence of “activated” RNA Pol II in osmotically stressed cells and Table 2, which demonstrates that relative transcript levels of many genes increase between 30 and 60 minutes, our data supports the hypothesis that RNA polymerase activity continues (perhaps more slowly) during acute hyperosmotic conditions.

## II. Translation

The GO term “ribosome” (GO:0005840) includes proteins involved in ribosome assembly/disassembly, localization and binding, as well as cytosolic and mitochondrial ribosome structure. Of the 205 mapped genes in this term, 86% have significantly increased transcript levels after 30 minutes of exposure to sucrose (Fig. 4a). Among the 10 genes with significantly decreased transcript levels are EIF2AK2, EIF2AK3, and EIF2AK4, which are kinases that phosphorylate translation initiation factor EIF2A to inhibit translation. Downregulation of these genes is consistent with increased translation. Transcript levels of all of the genes encoding the structural proteins of the cytosolic ribosome [18] increase by



**Figure 3.** Relative transcript levels of RNA Pol I, II and III protein subunit genes in HL-60/S4 cells exposed to tissue culture medium ( $\pm 300$  mM sucrose) for 30 and 60 minutes. Polymerase-specific subunit transcripts are displayed in individual panels, with shared subunit transcripts in a separate panel. See Table S5 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for complete data. Y-axis: Log<sub>2</sub>FC, log<sub>2</sub> of the ratio of transcript levels between the hyperosmotic stress and control conditions. Open bars signify that the change in transcript level is not significant (PPDE<0.95). Solid bars signify that the change in transcript level is significant (PPDE>0.95). Sucrose exposure times: 30 min (blue); 60 min (red). HGNC Gene codes are displayed above the relative transcript level bars.


a factor of about 2.5x and 3.3x after 30 and 60 minutes of exposure, respectively (Fig. 4b). In addition, of the 271 mapped genes in the term “ribosome biogenesis” (GO:0042254), 62% have significantly increased transcript levels after 30 minutes of exposure to sucrose (Fig. 4c). Given that it takes only a few minutes to translate the average-size protein [19, 20], we would expect that the cellular physiological state is changing continuously during the hyperosmotic stress.

## III. Mitochondria and Oxidative Phosphorylation

High concentrations of NaCl are known to cause depolarization of the mitochondrial membrane in a variety of cell types [3]. In our earlier study on the effects of acute sucrose stress upon undifferentiated HL-60/S4 cells [8], we determined that the mitochondrial membrane polarization is essentially normal for up to 1 hour of stress. Table 2 indicates that many GO terms associated with mitochondrial structure and function are enriched in genes with increased transcript levels, in cells exposed to sucrose (Fig. 5). These results suggest that acutely stressed cells are attempting to maintain ATP synthesis. It is of interest to note that desiccation tolerant plants and algae are reported to increase nuclear transcription of ribosomal protein genes [21-26] and mitochondrial protein genes [21,22], supporting a generalization of these responses in plants and animals.

## IV. Proteasome Activity

The GO term “proteasomal protein catabolic process” (GO:0010498) is enriched in both genes with increased




**Figure 4.** Relative transcript levels of ribosomal protein genes. For all panels, a dot represents one gene: a red dot indicates a significant change in transcript level (PPDE>0.95); a grey dot indicates a non-significant change (PPDE<0.95). Genes with log<sub>2</sub>FC values outside the range of the Y-axis are indicated by arrowheads. Figure 4a MA plots of all mapped genes associated with GO term “Ribosome” (GO:0005840) after 30 and 60 minutes of exposure to sucrose. The X-axis is the mean transcript level of exposed and unexposed (control) conditions. The Y-axis is log<sub>2</sub>FC of the ratio of transcript levels from exposed and unexposed (control) conditions. Figure 4b Scatterplots of transcript levels of structural protein genes of the cytosolic ribosome with linear regression lines demonstrating that despite relative differences in transcript levels across genes, all genes increase transcript levels by about 2.5x after 30 minutes and 3.3x after 60 minutes. Figure 4c MA plots of all mapped genes associated with GO term “Ribosome Biogenesis” (GO:0042254) after 30 and 60 minutes of exposure to sucrose.

and genes with decreased transcript levels (See Table S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>). Genes with increased transcript levels include nearly all those encoding the proteasome. In addition, ubiquitin genes UBB, UBC, UBA52, and RSP527A, as well as three E1 ubiquitin-activating enzymes, 13 E2 ubiquitin-conjugating enzymes, and components of the APC/C and ECS E3 ubiquitin ligase complexes have increased transcript levels. In contrast, genes encoding nine other E2 ubiquitin-conjugating enzymes and components of HCET, U-box, Cullin-Rbx, and single-RING finger E3 ubiquitin ligase complexes have decreased transcript levels. Together these results suggest an increase in proteasomal protein degradation mediated by increases and decreases in specific ubiquitination pathways.

#### V. Replication-Dependent histone mRNA

After 30 minutes of acute exposure to sucrose only 2% of genes have a log<sub>2</sub>FC>1 increase in transcript levels; by 60 minutes of exposure this increases to 4.5%. Among those transcript levels with a log<sub>2</sub>FC>1 increase, 15% encode rep-




**Figure 5.** Relative transcript levels of mitochondrial protein genes. All panels show MA plots where the X-axis is the mean transcript level of exposed and unexposed control conditions, and the Y-axis is log<sub>2</sub>FC of the ratio of transcript levels from exposed and unexposed control conditions. **Panel a**: Left, 30 minutes of exposure; Right, 60 minutes. Each dot represents one gene: red indicates a significant change in transcript level (PPDE>0.95); grey indicates a non-significant change (PPDE<0.95). (Panel a) Of the 230 genes in the Mitochondrial Protein Complex, 80% exhibit significantly increased relative transcript levels after 30 or 60 minutes of exposure. (Panel b) Of the 60 mapped genes in the Respiratory Chain Complex, 95% exhibit significantly increased relative transcript levels after 30 or 60 minutes of exposure. (Panel c) Of the 90 mapped genes in the Oxidoreductase Complex, 70% exhibit significantly increased relative transcript levels after 30 or 60 minutes of exposure.

lication-dependent (RD) histone genes. In fact, all the major classes of RD histones (H1, H2A, H2B, H3 and H4) showed significantly increased transcript levels in hyperosmotically-stressed HL-60/S4 cells (Fig. 6).

There are multiple copies (isoforms) for each of these histone classes arranged in clusters [27], with the largest cluster, HIST1, on chromosome 6 (6p21-6p22). A minor number of isoform copies are located on chromosome 1 in three clusters (HIST2, HIST3 and HIST4); only HIST1 and HIST2 clusters are significantly expressed in our data. Figure 6 displays the significant (PPDE>0.95) increased levels of histone class transcripts, combining RNA-seq reads for all isoforms for each histone gene class from both clusters HIST1 and HIST2. A complete listing of transcript level changes for the histone gene isoforms is shown in Table S6 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>.

The increase of RD histone transcript levels during hyperosmotic dehydration stress is surprising for two reasons: 1) As previously determined by cell cycle analyses [8], during this brief exposure to sucrose the % of cells in S phase was effectively unchanged (e.g., 0 min, 28.9%; 30 min, 29.9%; 60 min, 29.2%); 2) RD histone mRNAs normally have stem-loops, not 3' polyA tails [28,29]. Our method of mRNA purification selects for mRNAs with polyA tails. Why are these stressed and




**Figure 6.** Differential expression of polyA mRNA levels of the replication-dependent (RD) histone gene classes following acute hyperosmotic stress. Only isoforms with statistically significant changes (PPDE>0.95) were included for each gene class. Y-axis: Log<sub>2</sub>FC between the control and hyperosmotic stress condition for all isoforms of each histone class after 30 (blue) and 60 minutes (red). At 30 minutes of sucrose exposure, no histone H4 isoforms exhibited significantly increased transcript levels; at 60 minutes, one H4 isoform was significantly increased. See Table S6 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for a complete list of transcript level changes for the RD histone gene isoforms.

apparently quiescent cells showing an increase in RD histone mRNA transcript levels? One plausible explanation is that the histone locus body (HLB) is not functioning normally in hyperosmotically stressed cells, allowing RD histone mRNA with stem-loops to undergo a “default” conversion to 3’ polyA mRNA, resulting in an apparent increase in transcript levels. The function of many of these HLB component proteins are well understood [30-32]; for example, SLBP (Stem-Loop Binding Protein) is essential for stem-loop protection at all stages of histone mRNA metabolism and NPAT (Nuclear Protein, Coactivator of Histone Transcription) is a required scaffold for HLB formation. Transcript levels of SLBP and NPAT are significantly decreased during exposure to sucrose (see Table S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>), supporting the hypothesis that the HLBs are not structurally or functionally normal. Immunostaining for NPAT (Fig. 7) in cells fixed in sucrose and cells fixed in isosmotic buffer indicates that: **a)** HLBs are maintained during hyperosmotic stress, but are stained more weakly during stress; **b)** The number of HLBs per cell is lower in stressed cells. Unstressed (control) cells with functional HLBs convert only a small percentage of RD histone stem-loop RNA to 3’ polyA mRNA (see Table S6 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>, also [33]). This generates only trace amounts of RNA-Seq reads mapping to RD histone genes, compared to the stressed cells with presumptive “disabled” HLBs. Furthermore, evidence has been published that slow transcription (a conceivable consequence of hyperosmotic stress) prevents formation and stabilization of the stem-loop, leading to polyadenylation [32]. We suggest that the apparent increase in RD histone transcripts during hyperosmotic sucrose stress may represent a perturbation of the “normal” low-level stem-loop to polyA conversion, due to resultant changes in the composition and/or functioning of HLBs.

## VI. Reversing Chromatin Repression

Acute hyperosmotic stress appears to result in “relief” from transcription repression, as genes with decreased tran-




**Figure 7.** Immunostaining of NPAT (a major component of the HLB) before and after acute hyperosmotic stress for 30 minutes. Figure 7a Micrographs of HL-60/S4 cells stained with anti-NPAT (red dots) within interphase nuclei stained with DAPI (cyan). Note that the mitotic chromosomes (yellow arrowheads) are not stained. Note that the NPAT “dots” appear to be generally smaller and weaker stained in 300 mM sucrose, compared to 0 mM sucrose. Figure 7b Measured fraction of cells with discrete numbers of NPAT “dots” per cell in 0- or 300-mM sucrose. The “Mode” values of HLBs per cell are: 0 mM sucrose, 4 HLB dots/cell (blue); 300 mM sucrose, 2 HLB dots/cell (red).

script levels are over-represented in two GO terms involved in repression of transcription: “PcG (Polycomb Gene) Protein Complex” (GO:0031519), and “Transcription Repressor Complex” (GO:0017053). The Polycomb Gene Protein Complex includes three major groups, with complicated and overlapping interactomes [34] involved in epigenetic repression of transcription through histone modification. Key components of complexes in each group have decreased transcripts (Fig. S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>); of the 52 mapped genes associated with the GO term “Transcription Repressor Complex”, 27 experience a significant decrease in transcript levels following 30 minutes (28 genes after 60 minutes) exposure to sucrose. Collectively, these results suggest that transcription repressor effects are weakened or eliminated, consistent with the observed increase in transcript levels of many “building” genes.

## VII. Chromosome and chromatin structure

As documented in our earlier publication [8], protein components of the cohesin complex (e.g., RAD21 and CTCF) examined by immunostaining appear to separate (demix) from chromatin following exposure to 300 mM sucrose. This repositioning is expected to affect the structure of TADs (“Topologically Associated Domains”, closed chromatin loops consisting of self-interacting genomic regions [35, 36]), with possible effects upon gene expression. During examination of the transcriptome changes in hyperosmotic



**Figure 8.** Differential polyA mRNA levels of transcripts of chromatin structural protein genes following acute hyperosmotic stress. Y-axis: Log<sub>2</sub>FC between the control and hyperosmotic stress condition at 30 (blue) and 60 minutes (red). HGNC gene code names are displayed above the relative transcript level bars. The HGNC Name and the (Common Name) are presented here. **8a** Cohesin: ESPL1 (Separase), NIPBL (Cohesin Loading Factor). **8b** Condensin: SMC2 and SMC4 are subunits for both Condensin I and II. **8c** HMG and HP1: HMG1 (HMG14), HMG17 (HMG17), HMG2 (HMG2), CBX5 (HP1 $\alpha$ ), CBX1 (HP1 $\beta$ ), CBX3 (HP1 $\gamma$ ).

sucrose, we observed that many of the components of the cohesin “ring” structure exhibit significantly decreased transcript levels, relative to the transcriptome of unstressed cells (Fig. 8a). This observation argues that regulation of gene expression becomes increasingly aberrant during cell dehydration.

Condensin I and II also build “ring” structures around chromatin fibers [37,38]. Condensin I localizes on mitotic chromosomes; condensin II appears to be more important in organizing interphase chromatin. Loss of condensin can also affect gene expression, as well as centromere architecture. The actual consequences likely depend upon the chromatin context (e.g., whether the genes are active or inactive and what other proteins are bound). The consequences of changes in transcript levels in Condensin I or II to chromatin structure, during acute hyperosmotic stress, are not clear (Fig. 8b). Interphase and mitotic chromatin condensation [8] may be related to changing levels of condensins, but there is no evidence on this issue. SMC2 is a subunit component to both Condensin I and II. Immunostaining with anti-SMC2 demonstrates that it is part of an “axis” along the midline of mitotic chromosomes in normal isosmotic medium, but appears excluded from congealed mitotic chromosomes in 300 mM sucrose (See Fig. 5i and 5j in [8]).

Exploring relative transcript levels of various well-studied chromatin binding proteins (e.g., HMG and HP1) also indicates that their transcript levels change in response to acute hyperosmotic stress in directions consistent with decreased constraints on RNA Pol II transcription (Fig. 8c). Both groups of proteins can bind to nucleosomes and chromatin. To some extent, they have opposite functional consequences: HMG1 and HMG2 colocalize with epigenetic marks of active chromatin and with cell-type specific enhancers [39], counteracting H1 stabilization of chromatin higher-order structure [40]. HP1 proteins interact with histone tails and promote heterochromatin formation by phase separation [41,42]. The simultaneous increase of HMG1 protein transcript levels with a decrease in HP1 protein (CBX5) transcript levels argues for a relaxation of repression of gene expression during acute hyperosmotic stress.

**Discussion** Hyperosmotic stress in the human body is associated with inflammation and disease [2]. In addition,

many healthy tissues (e.g., kidneys, liver, lymphoid tissues, the cornea, gastrointestinal tract, intervertebral discs and cartilage in joints) experience transient hyperosmotic stress during normal functioning [2]. Blood cells circulating around and through these tissues should also experience acute osmotic changes. The renal (kidney) environment experiences very significant osmotic stress. For example, during the course of a normal night’s sleep (without water intake) urine can concentrate up to ~4x iso-osmotic “plasma” conditions [43]. Tissue-specific resilient responses allow cells to survive these osmotic stresses, but there have been few studies of genome-wide changes in gene transcription and, as far as we are aware, none involving sucrose and leukocytes.

The purpose of the present study was to gain some understanding of how the genome of an *in vivo* human cell responds to acute hyperosmotic (i.e., 30 and 60 minutes at ~600 milli-osmolar) stress. In our model cellular system, we exposed rapidly-growing “undifferentiated” human leukemic promyelocytes (HL-60/S4) to acute hyperosmotic stress (medium+300 mM sucrose, ~600 mOsM). These dehydrating conditions lead to rapid cell shrinkage, and interphase and mitotic chromatin condensation with apparent phase separation, including loss of colocalization (demixing) of various chromatin-associated proteins [8]. The cells remain alive and “healthy” for at least one hour, although longer exposure results in cell death within about 24 hours. Clearly complex biophysical changes are occurring within the shrinking cells, the crowding nuclei, and in the chemical environment surrounding nuclear chromatin [44]. We employed mRNA transcriptome analysis as an indicator of consequential genomic functional responses. Transcript levels at any time during the hyperosmotic stress represent the dynamic balance of mRNA transcription and degradation. It is of present interest that the transcript levels of major mRNA degradation factors and enzymes are significantly reduced (e.g., log<sub>2</sub>FC approximately -1 or lower) during the hyperosmotic stress: XRN1, the main mRNA 5’-3’ exonuclease; CNOT1, deadenylation; DCP2, decapping (See Table S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for the log<sub>2</sub>FC values after 30 and 60 minutes of dehydration.) The specific changes in mRNA transcript levels that we observe are difficult to explain in the face of an “apparent” general decrease in mRNA degradation factors.

Recent publications [45,46] address a complex “feedback” mechanism between mRNA degradation and transcription. Clearly, similar research approaches are needed in order to understand the dehydration effects upon transcript levels in the HL-60/S4 cell system.

Acute hyperosmotically-stressed undifferentiated HL-60/S4 cells exhibited differentially increased transcript levels for genes involved in transcription, translation, and mitochondrial function, as if the cells are embarking upon active growth. Increased transcript levels for proteasome functions also occurs, suggesting accelerated protein turnover. Perhaps the most surprising observation was the increase in transcript levels of replication-dependent (RD) histones. This was unexpected because the RD histone transcripts normally possess 3’stem-loops, rather than polyA tracks (which was the basis for the mRNA purification). Our speculation is that the histone locus body (HLB), which protects the stem-loops during S-Phase, is not functioning properly in the hyperosmotically-stressed cells. These presumptive “malfunctioning” HLBs may permit increased conversion of the stem-loops to 3’ polyA mRNA, producing an apparent increase in RD histone transcripts.

Genes with decreased transcript levels were enriched in various GO terms involved in repression of transcription: e.g. (GO:0031519) “PcG (Polycomb Gene) Protein Complex” and (GO:0017053) “Transcription Repressor Complex”), which might indicate a more permissive condition for mRNA synthesis (Fig. S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>). Other acute transcript level changes were seen in genes involved with 1) controlling chromatin domain structure (e.g., cohesin and condensin); 2) relaxing transcription repression and destabilizing heterochromatin (i.e., HMG and HP1 proteins).

Beyond the broad GO interpretations described above, the stressed cell physiology cannot be described as a stable state, especially in comparison to our previous analysis of chemically induced HL-60/S4 differentiation (for 4 days) into granulocytes and macrophage [13]. In that situation, the more stable phenotypic properties of the induced granulocyte and macrophage cell states were readily observed (microscopically) and in reasonable agreement with the corresponding transcriptome data. In contrast, the dehydrated HL-60/S4 transcriptome changes suggest (to us) a cellular attempt to build and grow, in the face of inevitable death. The future mode of cell death is not readily apparent in the transcriptomes. Examination for evidence of oncoming apoptosis is conflicting (e.g., transcript levels were significantly decreased for both “pro-apoptotic” Initiator and Effector caspases [CASP2 and CASP3] and for “anti-apoptotic” BCL2). Autophagy and necrosis also did not provide incriminating transcript changes. Nor is it evident that an “early osmotic stress response” [47] is occurring following the acute hyperosmotic stress. The relative transcript levels for early osmotic response genes NFAT5 (TonEBP) and SLC6A6 (TauT) both decrease sharply at 30 and 60 min. Immediate-Early Genes (IEG), such as the canonical FOS, JUN and MYC, can be transcribed in the first few minutes after cellular stress, initiating a rapid cellular response [48]. However, Fig. S2 at <https://postepybiochemii.ptbioch.edu.pl/>

[index.php/PB/](https://postepybiochemii.ptbioch.edu.pl/index.php/PB/) illustrates that for a well-established set of IEGs, the initial response of these genes is downregulation (with low statistical significance) in HL-60/S4 cells exposed to hyperosmotic sucrose for 30 and 60 min. Our observation is in contrast to an earlier study of isolated rat cardiomyocytes, which exhibited an increase in EGR1 and FOS mRNA following exposure to hyperosmotic conditions [49]. This suggests that the induction of IEG transcription by hyperosmotic stress may depend upon the type of cell.

One clue suggests that oxidative stress may play a role in the eventual cell death following hyperosmotic shock. Oxidative stress can occur when mitochondria produce more reactive oxygen species (ROS) than available antioxidant defenses can mitigate, resulting in critical cell damage. The transcript levels of many genes involved in glutathione- and thioredoxin-based antioxidant defense are elevated at 30 and 60 minutes of hyperosmotic stress, suggesting that ROS are increasing. In particular, the TXN2 product (thioredoxin 2) protects against oxidative stress in mitochondria, inducing cells to become insensitive to ROS-induced apoptosis [50]. If TXN2 transcript levels should decrease after 60 minutes exposure, or if the total amount of thioredoxin 2 (and other antioxidant proteins) are unable to compensate for the levels of ROS produced by hyperosmotic stress, ROS-induced apoptosis could be fatal [51].

Despite this lack of a complete understanding of the physiological state of the hyperosmotically-stressed cells, there is compelling evidence that the structure of chromatin is profoundly altered. This is readily seen in the earlier immunostaining study [8], which indicated an apparent phase separation and demixing of cohesin (RAD21) and condensin (SMC2) components, CTCF, histones H1.2 and H1.5, and non-histones HMGN2 and HMGB2. An independent study [52] employing hypertonic NaCl (which is also hyperosmotic) on breast cancer T47D cells analyzed chromatin changes using Hi-C and ChIP-seq (i.e., defining the DNA sites of bound RNA Pol II, CTCF and RAD21). Hypertonic stress resulted in a decrease in the number of TADs, a weakening of TAD boundaries and major perturbations of Compartments A and B (i.e., “euchromatin” and “heterochromatin”, respectively). RNA Pol II became dislocated from many transcription start sites (TSS) and appeared to run-off transcription end sites (TES). In addition, CTCF and RAD21 were displaced from their normal binding sites, consistent with our immunostained images [8]. Recent studies underscore the importance of normal cohesin function to higher-order interphase nuclear architecture [53].

The concept of Liquid-Liquid Phase Separation (LLPS) or “condensates”, where cellular macromolecules can be either concentrated into microscopic phases or dissolved in a cellular solution, depending upon their specific “critical concentration for phase separation”, has become a major biophysical perspective for interpreting non-membranous cellular particles, including chromatin [54-59]. Accepting that acute hyperosmotic stress with either NaCl [52] or sucrose [8,60] produces profound structural changes of *in vivo* chromatin (described as “altered” phase separation), we argue that this perturbed chromatin may yield somewhat disorganized transcriptomes and aberrant cell physiological

states. We suggest that, during acute dehydration, condensates may concentrate or disperse molecules in disordered or nonfunctional ways, dictated by the changing biophysical environment. Furthermore, we suggest that “unstressed” chromatin regions normally possess biophysical microheterogeneity of their “critical concentration for phase separation”. Most likely, this biophysical microheterogeneity is due to local differences in protein composition, post-translational modifications and physiologic state at the moments prior to the acute hyperosmotic shock.

Disorganized gene expression can have rapid effects. Transcription and translation of average size mRNA and protein (e.g., 500 amino acids) takes ~1 minute for each process [19,20]. The population of cellular proteins might exhibit significant changes by 30 and 60 minutes of osmotic stress. Decreases and/or displacement of chromatin structural proteins (e.g., components of cohesin and condensin) during acute osmotic shock could produce disruption of TADs [61], which may further exacerbate transcription disorder. From this perspective, we should not be surprised that cell differentiation, which follows evolutionary determined transcription programs in a “methodical” manner, yields coordinated transcriptomal responses; whereas, acute hyperosmotic stress and cell dehydration results in transcriptomes that are less focused on specific phenotypes. In the present study, the broad GO functions exhibiting relatively increased transcript levels (e.g., transcription, translation and mitochondrial oxidative phosphorylation) suggest that the “stressed” chromatin can still yield transcriptomes with some functional coordination. We suggest that this hyperosmotic stress response provides a useful *in vivo* model for examining rapid nuclear and chromatin biophysical changes (e.g., altered phase separation) that can influence chromatin higher-order structure and the regulation of transcription.

## CONCLUSION

The induced physiological state of acute hyperosmotically stressed undifferentiated HL-60/S4 cells resembles a rapid “attempt” to rebuild the damaged cells by increasing transcript levels of genes involved in transcription, translation, and energy production. The stressed cell physiologic state is not as well defined as the chemically-induced differentiated cell states of HL-60/S4, designated “granulocytes” and “macrophage” [13,14]. The suddenness of hyperosmotic stress with resultant changes in solute concentrations and macromolecular crowding, very likely yields “altered” phase separation with somewhat disordered gene expression, ultimately unable to prevent cell death.

## MATERIALS AND METHODS

### CELL CULTURE

HL-60/S4 cells (ATCC CRL-3306) were cultivated in RPMI-1640 medium, plus 10% FCS and 1% Pen/Strep. Rapidly growing cells in medium were added to dry sucrose in T-25 flasks yielding 300 mM sucrose (~600 mOsM, total) for 30 and 60 minutes, as described earlier [8]. Cell differentiation of HL-60/S4 into granulocytes and macrophage has also been described earlier [13].

## RNA PURIFICATION

Quadruplicate samples of undifferentiated HL-60/S4 cells ( $5 \times 10^6$  cells/sample) exposed to 0, 30, or 60 minutes of sucrose (total 12 samples) were centrifuged, rapidly frozen and stored in liquid nitrogen (LN<sub>2</sub>). Samples were thawed by the addition of the RLT lysis buffer from the Qiagen RNeasy Mini Kit and RNA purified according to the manufacturer’s protocol. Care was taken to maintain RNAase-free conditions with RNAase Zap. RNA was eluted with molecular biology grade water, frozen in LN<sub>2</sub> and shipped on dry ice to the Marshall University Genomics Core Facility. QC determinations (all 12 samples had a RIN score of 10), preparation of the libraries (Illumina mRNA library preparation Kit) and sequencing (Illumina HiSeq1500) was carried out at the Core Facility. The resulting sequences are available from the NCBI Sequence Read Archive as BioProject PRJNA686972. Data and analyses of undifferentiated, retinoic acid (RA)-treated and phorbol ester (TPA)-treated HL-60/S4 cells can be found in a previous publication [13].

## DATA ANALYSES AND PRESENTATION

We identified genes with significantly differential transcript levels following the RSEM-EBSeq workflow outlined at <http://deweylab.github.io/RSEM> and used the sequences and annotation of UCSC human genome v19 (hg19) from Illumina igenome: ([https://support.illumina.com/sequencing/sequencing\\_software/igenome.html](https://support.illumina.com/sequencing/sequencing_software/igenome.html)). Hg19 was chosen, instead of hg38, so that the hyperosmotic transcriptomes would be more comparable to the previous transcriptomes of undifferentiated and differentiated HL-60/S4 cells [13]. Furthermore, hg38 is a refinement of the assembly, including primarily alternatively spliced transcripts and a greater annotation of nonprotein-coding genes. It and the new HGNC names will recover more genes, but most of them are newly discovered and poorly annotated, so unlikely to have much impact on ORA and GO analyses. A bash script of the workflow is presented in Text S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/>. Briefly, we used bowtie2 v2.3.2 to map paired-end reads to transcripts extracted from the reference genome, and calculated transcript level values using RSEM v1.3.0. RSEM uses a maximum likelihood expectation-maximization algorithm to estimate the transcript levels of isoforms from RNA-Seq reads [62]. We then calculated the significance of relative expression differences using EBSeq v1.2.0 with the ng-vector option for isoform-level analysis [63]. EBSeq returns the normalized mean count of reads mapped using the median of ratios approach of DESeq [64] and the posterior probability of differential expression (PPDE) between control and treatment conditions, which is naturally corrected for multiple tests (i.e., the PPDE is equivalent to one minus the false discovery rate, FDR). We tested control versus 30 minutes and control *versus* 60 minutes of sucrose exposure, and accepted genes with PPDE>0.95 as having significantly different transcript levels. See Table S1 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for a complete table of EBseq normalized mean counts and PPDE values for each hg19 gene. Output from RSEM and EBSeq were loaded into a MySQL database with hg19 annotation for analysis. We uploaded lists of differentially expressed genes (i.e.,

HGNC gene codes) to WebGestalt (<http://bioinfo.vanderbilt.edu/webgestalt>) for (ORA) over-representation-analysis of GO non-redundant terms [65]. We considered a GO term over-represented if the hypergeometric test returned an adjusted  $p$  value (FDR) less than 0.05. See Tables S2, S3 and S4 at <https://postepybiochemii.ptbioch.edu.pl/index.php/PB/> for a detailed output for all GO terms significantly over-represented after 30 minutes or 60 minutes of exposure to sucrose.

## MICROSCOPY

STED imaging of DAPI stained HL-60/S4 cells  $\pm 300$  mM sucrose and Leica SP8 confocal imaging of HL-60/S4 cells  $\pm 300$  mM sucrose immunostained with rabbit anti-pPol II, rabbit anti-NPAT, mouse anti-epichromatin (PL2-6) and DAPI were performed by the same methods as described earlier [15]. Antibodies employed were HUABIO anti-Phospho RNA polymerase II POLR2A (S5) and Invitrogen (Thermo Fisher Scientific) anti-NPAT (PA5-668.39).

## REFERENCES

- Hancock R, Hadj-Sahraoui Y (2009) Isolation of cell nuclei using inert macromolecules to mimic the crowded cytoplasm. *PLoS One* 4(10): e7560
- Brocker C, Thompson DC, Vasiliou V (2012) The role of hyperosmotic stress in inflammation and disease. *Biomol Concepts* 3(4): 345-364
- Burg MB, Ferraris JD, Dmitrieva NI (2007) Cellular response to hyperosmotic stresses. *Physiol Rev* 87(4): 1441-74
- Richter K, Nessling M, Lichter P (2007) Experimental evidence for the influence of molecular crowding on nuclear architecture. *J Cell Sci* 120(Pt 9): 1673-80
- Finan JD, Guilak F (2010) The effects of osmotic stress on the structure and function of the cell nucleus. *J Cell Biochem* 109(3): 460-7
- Finan JD, Leddy HA, Guilak F (2011) Osmotic stress alters chromatin condensation and nucleocytoplasmic transport. *Biochem Biophys Res Commun* 408(2): 230-5
- Irianto J, Swift J, Martins RP, McPhail GD, Knight MM, Discher DE, Lee DA (2013) Osmotic challenge drives rapid and reversible chromatin condensation in chondrocytes. *Biophys J* 104(4): 759-69
- Olins AL, Gould TJ, Boyd L, Sarg B, Olins DE (2020) Hyperosmotic stress: in situ chromatin phase separation. *Nucleus* 11(1): 1-18
- Hoffmann EK, Lambert IH, Pedersen SF (2009) Physiology of cell volume regulation in vertebrates. *Physiol Rev* 89(1): 193-277
- Hosseiniyan Khatibi SM, Zununi Vahed F, Sharifi S, Ardalan M, Mohajel Shoja M, Zununi Vahed S (2019) Osmolytes resist against harsh osmolarity: Something old something new. *Biochimie* 158: 156-164
- Ortells MC, Morancho B, Drews-Elger K, Viollet B, Laderoute KR, López-Rodríguez C, Aramburu J (2012) Transcriptional regulation of gene expression during osmotic stress responses by the mammalian target of rapamycin. *Nucleic Acids Res* 40(10): 4368-84
- Arsenijevic T, Vujovic A, Libert F, Op de Beeck A, Hébrant A, Janssens S, Grégoire F, Lefort A, Bolaky N, Perret J, Caspers L, Willermain F, Delporte C (2013) Hyperosmotic stress induces cell cycle arrest in retinal pigmented epithelial cells. *Cell Death Dis* 4(6): e662
- Mark Welch DB, Jauch A, Langowski J, Olins AL, Olins DE (2017) Transcriptomes reflect the phenotypes of undifferentiated, granulocyte and macrophage forms of HL-60/S4 cells. *Nucleus* 8(2): 222-237
- Olins AL, Buendia B, Herrmann H, Lichter P, Olins DE (1998) Retinoic acid induction of nuclear envelope-limited chromatin sheets in HL-60. *Exp Cell Res* 245(1): 91-104
- Gould TJ, Tóth K, Mücke N, Langowski J, Hakusui AS, Olins AL, Olins DE (2017) Defining the epichromatin epitope. *Nucleus* 8(6): 625-640
- Zhou BR, Yadav KNS, Borgnia M, Hong J, Cao B, Olins AL, Olins DE, Bai Y, Zhang P (2019) Atomic resolution cryo-EM structure of a native-like CENP-A nucleosome aided by an antibody fragment. *Nat Commun* 10(1): 2301
- Cramer P (2019) Organization and regulation of gene transcription. *Nature* 573(7772): 45-54
- de la Cruz J, Karbstein K, Woolford Jr. JL (2015) Functions of ribosomal proteins in assembly of eukaryotic ribosomes in vivo. *Annu Rev Biochem* 84: 93-129
- Milo R, Phillips R (2015) *Cell Biology by the Numbers*, Garland Science, New York, USA
- Alberts BD, Lewis J, Raff M, Roberts K, Watson JD (1994) *Molecular Biology of The Cell*, Garland Publishing, New York, USA
- Oliver MJ, Dowd SE, Zaragoza J, Mauget SA, Payton PR (2004) The rehydration transcriptome of the desiccation-tolerant bryophyte *Tortula ruralis*: transcript classification and analysis. *BMC Genomics* 5: 89
- Peredo EL, Cardon ZG (2020) Shared up-regulation and contrasting down-regulation of gene expression distinguish desiccation-tolerant from intolerant green algae. *Proc Natl Acad Sci U S A* 117(29): 17438-17445
- Gao B, Li X, Zhang D, Liang Y, Yang H, Chen M, Zhang Y, Zhang J, Wood AJ (2017) Desiccation tolerance in bryophytes: The dehydration and rehydration transcriptomes in the desiccation-tolerant bryophyte *Bryum argenteum*. *Sci Rep* 7(1): 7571
- Gao B, Zhang D, Li X, Yang H, Zhang Y, Wood AJ (2015) De novo transcriptome characterization and gene expression profiling of the desiccation tolerant moss *Bryum argenteum* following rehydration. *BMC Genomics* 16(1): 416
- Gao B, Zhang D, Li X, Yang H, Wood AJ (2014) De novo assembly and characterization of the transcriptome in the desiccation-tolerant moss *Syntrichia caninervis*. *BMC Res Notes* 7: 490
- Yobi A, Schlauch KA, Tillett RL, Yim WC, Espinoza C, Wone BW, Cushman JC, Oliver MJ (2017) *Sporobolus stapfianus*: Insights into desiccation tolerance in the resurrection grasses from linking transcriptomics to metabolomics. *BMC Plant Biol* 17(1): 67
- Singh R, Bassett E, Chakravarti A, Parthun MR (2018) Replication-dependent histone isoforms: a new source of complexity in chromatin structure and function. *Nucleic Acids Res* 46(17): 8665-8678
- Marzluff WF, Wagner EJ, Duronio RJ (2008) Metabolism and regulation of canonical histone mRNAs: life without a poly(A) tail. *Nat Rev Genet* 9(11): 843-54
- Romeo V, Schümperli D (2016) Cycling in the nucleus: regulation of RNA 3' processing and nuclear organization of replication-dependent histone genes. *Curr Opin Cell Biol* 40: 23-31
- Duronio RJ, Marzluff WF (2017) Coordinating cell cycle-regulated histone gene expression through assembly and function of the Histone Locus Body. *RNA Biol* 14(6): 726-738
- Marzluff WF, Koreski KP (2017) Birth and Death of Histone mRNAs. *Trends Genet* 33(10): 745-759
- Saldi T, Fong N, Bentley DL (2018) Transcription elongation rate affects nascent histone pre-mRNA folding and 3' end processing. *Genes Dev* 32(3-4): 297-308
- Shepard PJ, Choi EA, Lu J, Flanagan LA, Hertel KJ, Shi Y (2011) Complex and dynamic landscape of RNA polyadenylation revealed by PAS-Seq. *RNA* 17(4): 761-72
- Hauri S, Comoglio F, Seimiya M, Gerstung M, Glatzer T, Hansen K, Aebersold R, Paro R, Gstaiger M, Beisel C (2016) A High-Density Map for Navigating the Human Polycomb Complexome. *Cell Rep* 17(2): 583-595
- Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragozcy M, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke A, Stamatoyannopoulos J, Mirny LA, Lander ES, Dekker J (2009) Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326(5950): 289-93
- McArthur E, Capra JA (2021) Topologically associating domain boundaries that are stable across diverse cell types are evolutionarily constrained and enriched for heritability. *Am J Hum Genet* 108(2): 269-283

37. Yuen KC, Gerton JL (2018) Taking cohesin and condensin in context. *PLoS Genet* 14(1): e1007118
38. Skibbens RV (2019) Condensins and cohesins - one of these things is not like the other! *J Cell Sci* 132(3): jcs220491
39. He B, Deng T, Zhu I, Furusawa T, Zhang S, Tang W, Postnikov Y, Ambs A, Li CC, Livak F, Landsman D, Bustin M (2018) Binding of HMGN proteins to cell specific enhancers stabilizes cell identity. *Nat Commun* 9(1): 5240
40. Murphy KJ, Cutter AR, Fang H, Postnikov YV, Bustin M, Hayes JJ (2017) HMGN1 and 2 remodel core and linker histone tail domains within chromatin. *Nucleic Acids Res* 45(17): 9917-9930
41. Sanulli S, Trnka MJ, Dharmarajan V, Tibble RW, Pascal BD, Burlingame AL, Griffin PR, Gross JD, Narlikar GJ (2019) HP1 reshapes nucleosome core to promote phase separation of heterochromatin. *Nature* 575(7782): 390-394
42. Sanulli S, Gross JD, Narlikar GJ (2019) Biophysical Properties of HP1-Mediated Heterochromatin. *Cold Spring Harb Symp Quant Biol* 84: 217-225
43. Sands JM, Layton HE (2009) The physiology of urinary concentration: an update. *Semin Nephrol* 29(3): 178-95
44. Hancock R (2014) The crowded nucleus. *Int Rev Cell Mol Biol* 307: 15-26
45. Hartenian E, Glaunsinger BA (2019) Feedback to the central dogma: cytoplasmic mRNA decay and transcription are interdependent processes. *Crit Rev Biochem Mol Biol* 54(4): 385-398
46. Chappleboim A, Joseph-Strauss D, Gershon O, Friedman N (2022) Transcription feedback dynamics in the wake of cytoplasmic mRNA degradation shutdown. *Nucleic Acids Res* 50(10): 5864-5880
47. Christoph K, Beck FX, Neuhofer W (2007) Osmoadaptation of Mammalian cells - an orchestrated network of protective genes. *Curr Genomics* 8(4): 209-18
48. Bahrami S, Drabløs F (2016) Gene regulation in the immediate-early response process. *Adv Biol Regul* 62: 37-49
49. Wollnik B, Kubisch C, Maass A, Vetter H, Neyses L (1993) Hyperosmotic stress induces immediate-early gene expression in ventricular adult cardiomyocytes. *Biochem Biophys Res Commun* 194(2): 642-6
50. Chen Y, Zhou Z, Min W (2018) Mitochondria, Oxidative Stress and Innate Immunity. *Front Physiol* 9: 1487
51. Redza-Dutordoir M, Averill-Bates DA (2016) Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim Biophys Acta* 1863(12): 2977-2992
52. Amat R, Böttcher R, Le Dily F, Vidal E, Quilez J, Cuartero Y, Beato M, de Nadal E, Posas F (2019) Rapid reversible changes in compartments and local chromatin organization revealed by hyperosmotic shock. *Genome Res* 29(1): 18-28
53. Cremer M, Brandstetter K, Maiser A, Rao SSP, Schmid VJ, Guirao-Ortiz M, Mitra N, Mamberti S, Klein KN, Gilbert DN, Leonhardt H, Cardoso MC, Aiden EL, Harz H, Cremer T (2020) Cohesin depleted cells rebuild functional nuclear compartments after endomitosis. *Nat Commun* 11(1): 6146
54. Banani SF, Lee HO, Hyman AA, Rosen MK (2017) Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* 18(5): 285-298
55. Shin Y, Brangwynne CP (2017) Liquid phase condensation in cell physiology and disease. *Science* 357(6357): eaaf4382
56. Shin Y, Chang YC, Lee DSW, Berry J, Sanders DW, Ronceray P, Wingreen NS, Haataja M, Brangwynne CP (2018) Liquid Nuclear Condensates Mechanically Sense and Restructure the Genome. *Cell* 175(6): 1481-1491
57. Larson AG, Narlikar GJ (2018) The Role of Phase Separation in Heterochromatin Formation, Function, and Regulation. *Biochemistry* 57(17): 2540-2548
58. Gibson BA, Doolittle LK, Schneider MWG, Jensen LE, Gamarra N, Henry L, Gerlich DW, Redding S, Rosen MK (2019) Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell* 179(2): 470-484
59. André AAM, Spruijt E (2020) Liquid-Liquid Phase Separation in Crowded Environments. *Int J Mol Sci* 21(16): 5908
60. Jaliha AP, ASchmidt A, Gao G, Little SR, Pitchiaya S, Walter NG (2021) Hyperosmotic phase separation: Condensates beyond inclusions, granules and organelles. *J Biol Chem* 296: 100044
61. Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kiefer-Kwon KR, Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J, Turner D, Ye Z, Omer AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander ES, Aiden EL (2017) Cohesin Loss Eliminates All Loop Domains. *Cell* 171(2): 305-320
62. Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12: 323
63. Leng N, Dawson JA, Thomson JA, Ruotti V, Rissman AI, Smits BM, Haag JD, Gould MN, Stewart RM, Kendziorski C (2013) EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics* 29(8): 1035-43
64. Anders S, Huber W (2010) Differential expression analysis for sequence count data. *Genome Biol* 11(10): R106
65. Zhang B, Kirov S, Snoddy J (2005) WebGestalt: an integrated system for exploring gene sets in various biological contexts. *Nucleic Acids Res* 33: W741-8

## SUPPORTING INFORMATION

Figure S1.

Figure S2.

Table S1 RSEM-EBSeq results.

Table S2 Bio

Table S3 Cell.

Table S4 Mol.

Table S5.

Table S6.

Text S1 BASH SCRIPT of the workflow.

## STATEMENTS & DECLARATIONS

**Funding.** DMW is supported by a gift from the G. Unger Vetlesen Foundation. TJG, ALO and DEO received funding from an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P20GM0103423. ALO and DEO received support from the School of Pharmacy, University of New England.

**Competing Interests.** The authors have no relevant financial or non-financial interests to disclose.

**Author Contributions.** DMW and DEO performed the data analyses. ALO performed the confocal microscopy. TJG contributed the STED images. ALO and DEO conceived the project, prepared the cells and mRNA. DMW, ALO and DEO wrote the paper. All authors read and approved the manuscript.

**Data Availability.** The RNA-seq data created in this study is openly available in the NCBI Sequence Read Archive as BioProject accession PRJNA686972.

**Ethics Approval.** No ethical approval is required since no animals or humans subjects were involved in the study.

**Consent to participate.** No human subjects participated in this study. Consent is not required.

**Consent to publish.** No human subjects participated in this study. Consent is not required.